Setting the **Global** Standard for Quality in Laboratory Testing

P: +1.610.688.0100 **Toll Free (US):** 877.447.1888 Suite 2500 F: +1.610.688.0700 E: customerservice@clsi.org

950 West Valley Road Wayne, PA 19087 USA



13 April 2018

To: Recipients of M60, 1st ed.

From: Jennifer K. Adams, MT(ASCP), MSHA

Vice President, Standards and Quality

Subject: Correction

This notification is to inform you of corrections made to CLSI document M60, Performance Standards for Antifungal Susceptibility Testing of Yeasts, 1st ed. The corrections are described below and shown as highlighted and/or stricken text in the table excerpt.

Table 5, Zone Diameter and Equivalent Minimal Inhibitory Concentration Breakpoints for Select Antifungal Agents Against *Candida* spp. After 24-Hour Incubation:

The resistant minimal inhibitory concentration (MIC) breakpoints for caspofungin and C. krusei and C. tropicalis are listed incorrectly as "> 1." The resistant MIC breakpoints were corrected to read "≥ 1."

Table 5. Zone Diameter and Equivalent Minimal Inhibitory Concentration Breakpoints for Select Antifungal Agents Against Candida spp. After 24-Hour Incubation

					Breakpoints and		Equivalent MIC Breakpoints and			
Antifungal		Interpretive Categories, mm*				Interpretive Categories, μg/mL				
Agent	Species	S	I	SDD	R	S	I	SDD	R	
Caspofungin	C. albicans	≥17	15–16	ı	≤14	≤0.25	0.5	_	≥1	
	C. glabrata	_	_	-	_	≤0.12	0.25	_	≥0.5	
	C. guilliermondii	≥13	11–12	ı	≤10	≤2	4	-	≥8	
	C. krusei	≥17	15–16	ı	≤14	≤0.25	0.5	_	<u>≥‡</u> ≥1	
	C. parapsilosis	≥13	11–12	-	≤10	≤2	4	_	≥8	
	C. tropicalis	≥17	15–16	ı	≤14	≤0.25	0.5	_	<u>≥4</u> ≥1	

If you require any additional clarification regarding these corrections, please contact CLSI Customer Service (customerservice@clsi.org).

We appreciate your commitment to CLSI and regret any inconvenience.