

Current Table 2F

- Currently includes many drugs not in use for the treatment of *N. gonorrhoeae*
- Suggest removal of
 - Cefoxitin
 - Cefuroxime
 - Cefmetazole
 - Cefotetan
 - Ceftazidime
 - Cefetamet

Table 2F. *Neisseria gonorrhoeae* (Continued)

Test/Report Group	Antimicrobial Agent	Disk Content
PENICILLINS		
O	Penicillin	10 units
CEPHEMS (PARENTERAL) (Including cephalosporins I, II,		
A	Ceftriaxone	30 µg
O	Cefoxitin	30 µg
O	Cefuroxime	30 µg
O	Cefepime	30 µg
O	Cefmetazole	30 µg
O	Cefotaxime	30 µg
O	Cefotetan	30 µg
O	Ceftazidime	30 µg
O	Ceftizoxime	30 µg
CEPHEMS (ORAL)		
A	Cefixime	5 µg
O	Cefpodoxime	10 µg
Inv.	Cefetamet	10 µg

Current Table 2F

- And removal of
 - Enofloxacin
 - Lomefloxacin
 - Ofloxacin
 - Fleroxacin

Table 2F. *Neisseria gonorrhoeae* (Continued)

Test/Report Group	Antimicrobial Agent	Disk Content
TETRACYCLINES		
(8) Organisms that are susceptible to tetracycline are also consid		
A	Tetracycline	30 µg
FLUOROQUINOLONES		
See general comment (3).		
A	Ciprofloxacin	5 µg
O	Enoxacin	10 µg
O	Lomefloxacin	10 µg
O	Ofloxacin	5 µg
Inv.	Fleroxacin	5 µg
AMINOCYCLITOLS		
O	Spectinomycin	100 µg

