

## Epidemiological Cutoff Values Eliminated From CLSI Document M100 Since 2010

Antimicrobial Agent	Disk Content	Zone Diameter ECV, mm		MIC ECV, µg/mL		M100 Edition in Which ECVs Were Last Included/Comments	Rationale
		WT	NWT	WT	NWT		
<b><i>Klebsiella (formerly Enterobacter) aerogenes, Enterobacter cloacae, Escherichia coli, Klebsiella pneumoniae, and Raoultella ornithinolytica</i></b>							
Colistin	-	-	-	≤ 2	≥ 4	M100, 29th ed.	MIC breakpoints were approved.
<b><i>Shigella spp.</i></b>							
Azithromycin	15 µg	≥ 16	≤ 15	≤ 8	≥ 16	M100, 30th ed.	For <i>Shigella flexneri</i> Disk diffusion and MIC breakpoints were approved for <i>Shigella spp.</i>
	-	-	-	≤ 16	≥ 32	M100, 30th ed.	For <i>Shigella sonnei</i> Disk diffusion and MIC breakpoints were approved for <i>Shigella spp.</i>
<b><i>Neisseria gonorrhoeae</i></b>							
Azithromycin	-	-	-	≤ 1	≥ 2	M100, 28th ed.	MIC breakpoints were approved.

Abbreviations: ECV, epidemiological cutoff value; MIC, minimal inhibitory concentration; NWT, non-wild-type; WT, wild-type.