

14 September 2021

To: Recipients of VET01S, 5th ed.  
 From: Jennifer K. Adams, MT(ASCP), MSHA  
 Vice President, Standards and Quality  
 Subject: Combined Corrections

This notice is intended to inform users of corrections made to CLSI document VET01S, *Performance Standards for Antimicrobial Disk and Dilution Susceptibility Tests for Bacteria Isolated From Animals*, 5th ed. The corrections are described below and shown as highlighted and/or stricken text in the table excerpts.

**Correction: 14 September 2021**

**Appendix D. Dosage Regimens Used to Establish Susceptible Veterinary-Specific Breakpoints:**

The Table 2F, *Bordetella bronchiseptica* tulathromycin dosage regimen for swine is listed incorrectly under Table 2G, *Mannheimia haemolytica*. The dosage regimen for swine has been correctly listed under Table 2F as “2.5 mg/kg IM once.”

The Table 2G *Mannheimia haemolytica* enrofloxacin dosage regimen for cattle is listed incorrectly as “7.5 mg/kg IM or SC once.” The dosage regimen for cattle has been corrected to read “7.5 mg/kg SC once.”

The Table 2G, *Mannheimia haemolytica* tulathromycin dosage regimen for cattle is listed incorrectly as “2.5 mg/kg IM once.” The dosage regimen for cattle has been corrected to read “2.5 mg/kg SC once.”

**Appendix D. Dosage Regimens Used to Establish Susceptible Veterinary-Specific Breakpoints**

Antimicrobial Agent	Breakpoints and Interpretive Categories			Comments
	MIC	Body Site	Dosage Regimen	
<b>Table 2F. <i>Bordetella bronchiseptica</i></b>				
Swine				
Tulathromycin	≤ 16	Resp	2.5 mg/kg IM once	
<b>Table 2G. <i>Mannheimia haemolytica</i></b>				
Cattle				
Tulathromycin	≤ 16	Resp	2.5 mg/kg IM once	
Enrofloxacin	≤ 0.25	Resp	7.5 mg/kg <del>IM or SC</del> once	See footnote b.
Tulathromycin	≤ 16	Resp	2.5 mg/kg <del>IM</del> SC once	

Correction: 30 June 2021

**Appendix D. Dosage Regimens Used to Establish Susceptible Veterinary-Specific Breakpoints:**

The Table 2G, *Mannheimia haemolytica* penicillin G dosage regimen for cattle is listed incorrectly as “33 000 U/kg IM by needle in neck every 24 hours.” The dosage regimen for cattle has been corrected to read “22 000 U/kg IM by needle in neck every 24 hours.”

**Appendix D. Dosage Regimens Used to Establish Susceptible Veterinary-Specific Breakpoints**

Antimicrobial Agent	Breakpoints and Interpretive Categories			Comments
	Susceptible			
	MIC	Body Site	Dosage Regimen	
Table 2G. <i>Mannheimia haemolytica</i>				
Cattle				
Penicillin G	≤0.25	Resp	<del>33 000</del> 22 000 U/kg IM by needle in neck every 24 hours	See footnote a.

If you require any additional clarification regarding these corrections, please contact CLSI Customer Service ([customerservice@clsi.org](mailto:customerservice@clsi.org)).

We appreciate your commitment to CLSI and regret any inconvenience.